

## Linear Motion Under a Variable Force 2



Q1.

A particle  $P$  of mass  $0.4 \text{ kg}$  is released from rest at the top of a smooth plane inclined at  $30^\circ$  to the horizontal. The motion of  $P$  down the slope is opposed by a force of magnitude  $0.6x \text{ N}$ , where  $x \text{ m}$  is the distance  $P$  has travelled down the slope.  $P$  comes to rest before reaching the foot of the slope. Calculate

- (i) the greatest speed of  $P$  during its motion, [7]
  - (ii) the distance travelled by  $P$  during its motion. [2]
- 

Q2.

A particle  $P$  of mass  $0.25 \text{ kg}$  moves in a straight line on a smooth horizontal surface. At time  $t \text{ s}$  the velocity of  $P$  is  $v \text{ m s}^{-1}$ . A variable force of magnitude  $3t \text{ N}$  opposes the motion of  $P$ .

- (i) Given that  $P$  comes to rest when  $t = 3$ , find  $v$  when  $t = 0$ . [4]
  - (ii) Calculate the distance travelled by  $P$  in the interval  $0 \leq t \leq 3$ . [3]
- 

Q3.

A particle  $P$  of mass  $0.2 \text{ kg}$  is projected horizontally from a fixed point  $O$ , and moves in a straight line on a smooth horizontal surface. A force of magnitude  $0.4x \text{ N}$  acts on  $P$  in the direction  $PO$ , where  $x \text{ m}$  is the displacement of  $P$  from  $O$ .

- (i) Given that  $P$  comes to instantaneous rest when  $x = 2.5$ , find the initial kinetic energy of  $P$ . [4]
  - (ii) Find the value of  $x$  on the first occasion when the speed of  $P$  is  $2 \text{ m s}^{-1}$ . [2]
- 

Q4.

A particle  $P$  of mass  $0.2 \text{ kg}$  is released from rest and falls vertically. At time  $t \text{ s}$  after release  $P$  has speed  $v \text{ m s}^{-1}$ . A resisting force of magnitude  $0.8v \text{ N}$  acts on  $P$ .

- (i) Show that the acceleration of  $P$  is  $(10 - 4v) \text{ m s}^{-2}$ . [2]
  - (ii) Find the value of  $v$  when  $t = 0.6$ . [5]
-

# Linear Motion Under a Variable Force 2

Q5.



Two particles  $P$  and  $Q$ , of masses  $0.4 \text{ kg}$  and  $0.2 \text{ kg}$  respectively, are attached to opposite ends of a light inextensible string.  $P$  is placed on a horizontal table and the string passes over a small smooth pulley at the edge of the table. The string is taut and the part of the string attached to  $Q$  is vertical (see diagram). The coefficient of friction between  $P$  and the table is  $0.5$ .  $Q$  is projected vertically downwards with speed  $5 \text{ m s}^{-1}$ , and at time  $t \text{ s}$  after the instant of projection the speed of the particles is  $v \text{ m s}^{-1}$ . The motion of each particle is opposed by a resisting force of magnitude  $0.9v \text{ N}$ . The particle  $P$  does not reach the pulley.

(i) Show that  $\frac{dv}{dt} = -3v$ . [4]

(ii) Find the value of  $t$  when the particles have speed  $2.5 \text{ m s}^{-1}$  and the distance that each particle has travelled in this time. [7]

Q6.

A particle of mass  $0.2 \text{ kg}$  is projected vertically downwards with initial speed  $4 \text{ m s}^{-1}$ . A resisting force of magnitude  $0.09v \text{ N}$  acts vertically upwards on the particle during its descent, where  $v \text{ m s}^{-1}$  is the downwards velocity of the particle at time  $t \text{ s}$  after being set in motion.

(i) Show that the acceleration of the particle is  $(10 - 0.45v) \text{ m s}^{-2}$ . [1]

(ii) Find  $v$  when  $t = 1.5$ . [5]

Q7.

A particle  $P$  of mass  $0.5 \text{ kg}$  moves in a straight line on a smooth horizontal surface. The velocity of  $P$  is  $v \text{ m s}^{-1}$  when the displacement of  $P$  from  $O$  is  $x \text{ m}$ . A single horizontal force of magnitude  $0.16e^x \text{ N}$  acts on  $P$  in the direction  $OP$ . The velocity of  $P$  when it is at  $O$  is  $0.8 \text{ m s}^{-1}$ .

(i) Show that  $v = 0.8e^{\frac{1}{2}x}$ . [6]

(ii) Find the time taken by  $P$  to travel  $1.4 \text{ m}$  from  $O$ . [4]

## Linear Motion Under a Variable Force 2



Q8.

A particle  $P$  of mass  $0.8 \text{ kg}$  moves along the  $x$ -axis on a horizontal surface. When the displacement of  $P$  from the origin  $O$  is  $x \text{ m}$  the velocity of  $P$  is  $v \text{ m s}^{-1}$  in the positive  $x$ -direction. Two horizontal forces act on  $P$ . One force has magnitude  $4e^{-x} \text{ N}$  and acts in the positive  $x$ -direction. The other force has magnitude  $2.4x^2 \text{ N}$  and acts in the negative  $x$ -direction.

(i) Show that  $v \frac{dv}{dx} = 5e^{-x} - 3x^2$ . [2]

(ii) The velocity of  $P$  as it passes through  $O$  is  $6 \text{ m s}^{-1}$ . Find the velocity of  $P$  when  $x = 2$ . [5]

---

Q9.

A small block  $B$  of mass  $0.2 \text{ kg}$  is placed at a fixed point  $O$  on a smooth horizontal surface. A horizontal force of magnitude  $0.42 \text{ N}$  is applied to  $B$ . At time  $t \text{ s}$  after the force is first applied, the velocity of  $B$  away from  $O$  is  $v \text{ m s}^{-1}$ .

(i) Find the value of  $v$  when  $t = 1$ . [2]

For  $t > 1$  an additional force, of magnitude  $0.32t \text{ N}$  and directed towards  $O$ , is applied to  $B$ . The force of magnitude  $0.42 \text{ N}$  continues to act as before.

(ii) Find the value of  $v$  when  $t = 2$ . [3]

For  $t > 2$  a third force, of magnitude  $0.06t^2 \text{ N}$  and directed away from  $O$ , is applied to  $B$ . The other two forces continue to act as before.

(iii) Show that the velocity of  $B$  is the same when  $t = 2$  and when  $t = 3$ . [3]

---